

**ehp**

**Environmental Health**

ehponline.org

P E R S P E C T I V E S

Published by the National Institute of  
Environmental Health Sciences

# Public Health and Economic Consequences of Methylmercury Toxicity to the Developing Brain

---

Leonardo Trasande, Philip J. Landrigan, Clyde Schechter  
doi:10.1289/ehp.7743 (available at <http://dx.doi.org/>)  
Online 28 February 2005



The National Institute of Environmental Health Sciences  
National Institutes of Health  
U.S. Department of Health and Human Services

## **Public Health and Economic Consequences of Methylmercury Toxicity to the Developing Brain**

Leonardo Trasande<sup>1,2,3,4</sup>  
Philip J. Landrigan<sup>1,2</sup>  
Clyde Schechter<sup>5</sup>

Corresponding Author: Leonardo Trasande, MD, MPP, Center for Children's Health and the Environment, Department of Community and Preventive Medicine, Mount Sinai School of Medicine, One Gustave L. Levy Place, Box 1057, New York, NY 10029; telephone: 212-241-8029; fax: 212-996-0407; email: [leo.trasande@mssm.edu](mailto:leo.trasande@mssm.edu)

<sup>1</sup>Center for Children's Health and the Environment, Department of Community and Preventive Medicine, New York, NY

<sup>2</sup>Department of Pediatrics, Mount Sinai School of Medicine, New York, NY

<sup>3</sup>Division of General Pediatrics, Children's Hospital, Boston, MA

<sup>4</sup>Department of Pediatrics, Harvard Medical School, Boston, MA

<sup>5</sup>Department of Family Medicine, Albert Einstein College of Medicine, Bronx, NY

**Running Title:** Cost of MeHg Toxicity to the Developing Brain

**Key Words:** Methylmercury, mercury, cord blood, lost economic productivity, power plants, electrical generation facilities, environmentally attributable fraction, fetal exposure, children's health, cognitive development

**Grant Information:** This research was supported by a grant from the National Institute of Environmental Health Sciences (NIEHS) through the Superfund Basic Research Program (P42 ES07384 to Mount Sinai School of Medicine), and by the Jenifer Altman Foundation, Physicians for Social Responsibility and the Rena Shulsky Foundation.

**Acknowledgements:** We are grateful for the helpful advice of Drs. Philippe Grandjean, Esben Budtz-Jorgensen, Danielle Laraque, Paul Leigh and Judith Palfrey as well as to Drs. Dorothy P. Rice and Wendy Max for their assistance with lifetime earnings estimates.

The authors have no conflicts of interest to report.

Abstract Word Count 235

Article Word Count 5720 + 3 figures (750) = 6470



## **Abbreviations**

µg/L: micrograms per Liter

BMDL: Benchmark dose level

DHA: docosahexaenoic acid

EAF: Environmentally attributable fraction

EPA: Environmental Protection Agency

IOM: Institute of Medicine

IQ: Intellectual Quotient

LCPUFAs: Long-chain polyunsaturated fatty acids

NAS: National Academy of Sciences

NHANES: National Health and Nutrition Examination Survey

WISC-R: Wechsler Intelligence Scale-Revised

## **Outline of MS**

Abstract

Introduction

Methodology

Environmentally attributable fraction (EAF) model

Toxic Effects of Methylmercury Exposure

Impact of Methylmercury Exposure on IQ

Calculation of Economic Costs of IQ Loss

American Sources of Mercury Emission

Results

Discussion

## **Abstract**

Methylmercury is a developmental neurotoxicant. Exposure results principally from consumption by pregnant women of seafood contaminated by mercury from anthropogenic (70%) and natural (30%) sources. Throughout the 1990s, the Environmental Protection Agency made steady progress in reducing mercury emissions from anthropogenic sources, especially from power plants, which account for 41% of anthropogenic emissions. However, EPA recently proposed to slow this progress, citing high costs of pollution abatement. To put into perspective the costs of controlling emissions from American power plants, we have estimated the economic costs of methylmercury toxicity attributable to mercury from these plants. We used an environmentally attributable fraction model, and limited our analysis to the neurodevelopmental impacts, specifically loss of intelligence. Using national blood mercury prevalence data from the Centers for Disease Control and Prevention, we found that between 316,588 and 637,233 children each year have cord blood Hg levels  $>5.8 \mu\text{g/L}$ , a level associated with loss of IQ. The resulting loss of intelligence causes diminished economic productivity that persists over the entire lifetime of these children. This lost productivity is the major cost of methylmercury toxicity, and it amounts to \$8.7 billion annually (range: \$2.2-\$43.8 billion, 2000 dollars). Of this total, \$1.3 billion (range: \$0.1-\$6.5 billion) each year is attributable to mercury emissions from American power plants. This significant toll threatens the economic health and security of the United States and should be considered in the debate on mercury pollution controls.

## **Introduction**

Mercury is an ubiquitous environmental toxicant (Goldman et al. 2001). It exists in three forms, each of which possesses different bioavailability and toxicity: the metallic element, inorganic salts, and organic compounds (methylmercury, ethylmercury and phenylmercury). (Franzblau 1994) Although volcanoes and other natural sources release some elemental mercury to the environment, anthropogenic emissions from coal-fired electric power generation facilities, chloralkali production, waste incineration and other industrial activities now account for approximately 70% of the 5500 metric tons of mercury that are released into the earth's atmosphere each year (United Nations Environmental Programme 2002). Elemental mercury is readily aerosolized due to its low boiling point, and once airborne it can travel long distances to eventually deposit into soil and water. In the sediments of rivers, lakes and the ocean, metallic mercury is transformed within microorganisms into methylmercury (Guiamares et al. 2000). This methylmercury biomagnifies in the marine food chain to reach very high concentrations in predatory fish such as swordfish, tuna, king mackerel and shark (Dietz et al. 2000; Gilmour et al 2000; Mason et al. 1995; Neumann et al. 1999). Consumption of contaminated fish is the major route of human exposure to methylmercury.

The toxicity of methylmercury to the developing brain was first recognized in the 1950s in Minamata, Japan, where consumption of fish with high concentrations of methylmercury by pregnant women resulted in at least thirty cases of cerebral palsy in children; exposed women were affected minimally if at all (Harada 1968). A similar episode followed in 1972 in Iraq when the use of a methylmercury fungicide led to poisoning in thousands of people (Bakir

1973); again infants and children were most profoundly affected (Amin-Zaki et al. 1974; Amin-Zaki et al. 1979). The vulnerability of the developing brain to methylmercury reflects the ability of lipophilic methylmercury to cross the placenta and concentrate in the central nervous system. (Campbell et al. 1992) Moreover, the blood-brain barrier is not fully developed until after the first year of life, and methylmercury can cross this incomplete barrier. (Rodier 1995)

Three recent, large-scale prospective epidemiologic studies have examined children who experienced methylmercury exposures *in utero* at concentrations relevant to current US exposure levels. The first of these studies, a cohort in New Zealand, found a three-point decrement in the Wechsler Intelligence Scale-Revised (WISC-R) full-scale IQ among children born to women with maternal hair mercury concentrations above 6 µg/g (Kjellstrom et al. 1986; Kjellstrom et al. 1989). A second study in the Seychelles Islands in the Indian Ocean found only one adverse association with maternal hair mercury concentration among forty-eight neurodevelopmental endpoints examined (prolonged time to complete a grooved pegboard test with the nonpreferred hand) (Myers et al. 2003). However, the grooved pegboard test was one of the few neurobehavioral instruments in the Seychelles study not subject to the vagaries of translation that can degrade the validity of culture-bound tests of higher cognitive function when they are applied in developing nations (Landrigan and Goldman 2003). A third prospective study in the Faroe Islands, a component of Denmark inhabited by a Scandinavian population in the North Atlantic, has followed a cohort of children for fourteen years, and collected data on seventeen neurodevelopmental endpoints, as well as on the impact of methylmercury on cardiovascular function. The Faroes researchers found significant dose-related, adverse associations between prenatal mercury exposure and performance on a wide range of memory, attention, language and

visual-spatial perception tests (Grandjean et al. 1997). The significance of these associations remained evident when blood levels of polychlorinated biphenyls, which are known developmental neurotoxicants, (Jacobson and Jacobson 1996) were included in the analysis (Budtz-Jorgensen et al. 2002; Steuerwald et al., 2000). Methylmercury exposure was also associated with decreased sympathetic- and parasympathetic-mediated modulation of heart rate variability (Grandjean et al 2004) and with persistent delays in peak I-III brainstem evoked potentials (Murata et al 2004).

An assessment of these three prospective studies by the National Academy of Sciences (NAS) concluded that there is strong evidence for the fetal neurotoxicity of methylmercury, even at low concentrations of exposure. Moreover, the NAS opined that the most credible of the three prospective epidemiologic studies was the Faroe Islands investigation. In recommending a procedure for setting a reference dose for a methylmercury standard, the NAS chose to use a linear model to represent the relationship between mercury exposure and neurodevelopmental outcomes, and based this model on the Faroe Islands data. The NAS found that the cord blood methylmercury concentration was the most sensitive biomarker of exposure *in utero* and correlated best with neurobehavioral outcomes. The NAS was not deterred by the apparently negative findings of the Seychelles Islands study, which it noted was based on a smaller cohort than the Faroe Islands investigation and had only 50% statistical power to detect the effects observed in the Faroes (NRC 2000).

Since January 2003, the issue of early life exposure to methylmercury has become the topic of intense debate after the Environmental Protection Agency (EPA) announced a proposal to

reverse strict controls on emissions of mercury from coal-fired power plants. This proposed “Clear Skies Act” would slow recent progress in controlling mercury emission rates from electric generation facilities and would allow these releases to remain as high as twenty-six tons per year through 2010 (U.S. EPA 2004a). By contrast, existing protections under the Clean Air Act will limit mercury emissions from coal-fired power plants to five tons per year by 2008 (U.S. EPA 2004b). EPA’s technical analyses in support of “Clear Skies” failed to incorporate or quantify consideration of the health impacts resulting from increased mercury emissions (U.S. EPA 2004c). After legislative momentum for this proposal faded, EPA proposed an almost identical Utility Mercury Reductions Rule which again failed to examine impacts on health. The EPA is actively reviewing comments in preparation for issuing a final Rule in March 2005 (U.S. EPA 2004d).

To assess the costs that may result from exposure of the developing brain to methylmercury, we estimated the economic impact of anthropogenic methylmercury exposure in the 2000 U.S. birth cohort. We calculated the fraction of this cost that could be attributed to mercury emitted by American electric power generation facilities.

## **Methodology**

*Environmentally attributable fraction (EAF) model.* To assess the disease burden and the costs due to methylmercury exposure, we used an environmentally attributable fraction (EAF) model. The EAF approach was developed by the Institute of Medicine (IOM) to assess the "fractional contribution" of the environment to causation of illness in the U.S. (IOM 1981), and it has been

used to assess the costs of environmental and occupational disease (Fahs et al. 1989; Leigh et al. 1997). It was used recently to estimate the environmentally attributable costs of lead poisoning, asthma, pediatric cancer, and neurodevelopmental disabilities in American children (Landrigan et al. 2002). The EAF is defined by Smith *et al.* as "the percentage of a particular disease category that would be eliminated if environmental risk factors were reduced to their lowest feasible concentrations" (Smith et al. 1999). The EAF is a composite value and is the product of the prevalence of a risk factor multiplied by the relative risk of disease associated with that risk factor. Its calculation is a useful tool in developing strategies for resource allocation and prioritization in public health. The general model developed by the IOM and used in the present analysis is the following:

$$\text{Costs} = \text{Disease rate} \times \text{EAF} \times \text{Population size} \times \text{Cost per case}$$

In this equation, EAF represents the environmentally attributable fraction. "Cost per case" refers to discounted lifetime expenditures attributable to a particular disease including direct costs of health care, costs of rehabilitation, and lost productivity. The terms "disease rate" and "population size" refer, respectively, to either the incidence or prevalence of a disease and the size of the population at risk.

In applying the EAF model, we first reviewed the adverse effects of methylmercury exposure. We then estimated the costs of those effects, and subsequently applied a further fraction to parse out the cost of anthropogenic methylmercury exposure resulting from emissions of American electrical generation facilities.

*Toxic Effects of Methylmercury Exposure.* The NAS found neurodevelopmental effects in the children of women who had consumed fish and seafood during pregnancy to be the most important and best-studied endpoint for methylmercury toxicity. Though the NAS identified other potentially significant toxicities resulting from methylmercury exposure, such as nephrotoxicity and carcinogenicity, those effects were less well characterized (NRC 2000). We therefore chose to limit our analysis to the neurodevelopmental impact of methylmercury toxicity.

There is no evidence to date validating the existence of a threshold blood mercury concentration below which adverse effects on cognition are not seen. EPA has, however, set a benchmark cord blood mercury dose concentration (BMDL) of 58  $\mu\text{g/L}$ . This level that corresponds to the lower limit of the 95% confidence interval for the concentration at which there is a doubling in the Faroes study in the prevalence of test scores (5% to 10%) in the clinically subnormal range for the Boston Naming Test (Rice et al. 2003). It is important to note that this is not a concentration below which no observed adverse effects were found. The Faroes and New Zealand cohorts both support the conclusion that developmental effects become apparent at levels of approximately one part per million mercury in hair, or 5.8  $\mu\text{g/L}$  in cord blood (Grandjean et al. 1997; Kjellstrom et al. 1986; Kjellstrom et al. 1989). The Faroes study also found that effects on delayed brainstem auditory responses occurred at much lower exposure concentrations (Murata et al. 2004). In its report, the National Academy of Sciences (NAS) concluded that the likelihood of subnormal scores on neurodevelopmental tests following *in utero* exposure to methylmercury increased as cord blood concentrations increased from levels as low as 5  $\mu\text{g/L}$  to the BMDL of

58 µg/L (NRC 2000). In light of those findings, we made the decision in this analysis to apply a no adverse effect level of 5.8 µg/L, the lowest level at which adverse neurodevelopmental effects were demonstrated in the cohort studies.

Recent data suggest that the cord blood mercury concentration may on average be 70% higher than the maternal blood mercury concentration (Stern and Smith 2003), and a recent analysis suggests that a modification of the EPA reference dose for methylmercury be made to reflect a cord blood/maternal blood ratio that is greater than 1 (Stern 2005). If the developmental effects of mercury exposure do, in fact, begin at 5.8 µg/L in cord blood, as suggested by the Faroes (Grandjean et al. 1997) and New Zealand (Kjellstrom et al. 1986; Kjellstrom et al. 1989) data and by the NAS report (NRC 2000), then effects would occur in children born to women of child-bearing age with blood mercury concentrations  $\geq 3.41$  ( $=5.8/1.7$ ) µg/L. National population data from the 1999-2000 National Health and Nutrition Examination Survey (NHANES) found that 15.7% of American women of childbearing age have total blood mercury concentrations  $\geq 3.5$  µg/L. (Mahaffey et al. 2004)

To compute IQ decrements in infants that have resulted from these elevated maternal mercury exposures, we used published data on percentages of women of childbearing age with mercury concentrations at or above 3.5, 4.84, 5.8, 7.13 and 15.0 µg/L. We assumed conservatively that all mercury concentrations within each of the segments of the distribution were at the lower bound of the range. We assumed that the probability of giving birth to a child did not correlate with mercury level in a woman of childbearing age. In our base case analysis, we calculated economic costs assuming that children born to women with mercury concentrations 3.5-4.84

µg/L suffer no loss in cognition, and that successive portions of the birth cohort experience loss of cognition associated with cord blood levels of 8.2, 9.9, 12.1 and 25.5 µg/L, respectively.

Recently, the Faroes researchers reviewed their cohort data and found fetal blood mercury concentrations to be only 30% higher than maternal blood concentrations. (Budtz-Jorgensen et al. 2004) In light of these findings, and to avoid overestimation of the magnitude of impacts, we chose not to include children born to mothers with blood mercury concentrations between 3.5-4.84 µg/L in our base case analysis.

To assess the impact on our findings of a range of various possible ratios between maternal and cord blood mercury concentrations, we conducted a sensitivity analysis. In this analysis, we set as a lower bound for our estimate the costs to children with estimated cord blood concentrations  $\geq 5.8$  µg/L (assuming a cord/maternal blood ratio of 1), and assumed no IQ impact  $< 4.84$  µg/L (assuming a cord/maternal blood ratio of 1.19). This estimate assumed no loss of cognition to children born to women with mercury concentration  $< 5.8$  µg/L, and assumed that subsequent portions of the birth cohort experienced cord blood mercury concentrations of 5.8, 7.13 and 15 µg/L, respectively. To estimate economic costs in this scenario, we calculated no costs for children with blood mercury concentrations below 4.84 µg/L. We calculated costs resulting from an incremental increase in blood mercury concentration from 4.84 to 5.8 µg/L in the percentage of the population with blood mercury levels between 5.8 and 7.13 µg/L, and added those costs to the costs resulting from increases from 4.84 to 7.13 µg/L and 4.84 to 15 µg/L in the percentages of the population between 7.13 and 15 µg/L, and above 15 µg/L, respectively.

The result of this calculation is expressed in our analysis as a lower bound for the true economic cost of methylmercury toxicity to the developing brain.

*Impact of Methylmercury Exposure on IQ.* The Faroes study found that a doubling of mercury concentration was associated with adverse impacts on neurodevelopmental tests ranging from 5.69-15.93% of a standard deviation (Grandjean et al. 1999). Assuming that IQ is normally distributed with a standard deviation of 15 points, a doubling of mercury concentration would be associated with a decrement ranging from 0.85-2.4 IQ points. The Faroes researchers utilized a structural equation analysis to produce estimates of impact of methylmercury on verbal and motor function at seven years of age, and found an association between a doubling of blood mercury and loss of 9.74% of a standard deviation on motor function and of 10.45% a standard deviation on verbal function (Budtz-Jorgensen et al. 2002). This analysis suggests that a doubling in mercury concentration produces a decrement of approximately 10% of a standard deviation, or 1.5 IQ points. In the New Zealand study, the average WISC-R full-scale IQ for the study population (n=237) was 93. In the group with maternal hair mercury above 6 µg/g (about 4-fold higher than in the study population, n=61) the average was 90 (Kjellstrom et al. 1999). This finding provides further support for our use of a loss of 1.5 IQ points for each doubling in our base case analysis. Confounders such as polychlorinated biphenyls were found not to cause significant confounding of the data in the Faroe Islands study (Budtz-Jorgensen et al. 2002; Steuerwald et al. 2000). As a conservative measure, we nonetheless chose to set as outer bounds for the impact on intelligence of methylmercury exposure a range of IQ decrements from 0.85-2.4 IQ points per doubling, as described by the Faroes researchers (Jorgensen 2004). In applying

the EAF methodology, we assume that the relationship between cord blood mercury and IQ is relatively linear over the range of exposures studied ( $>5.8 \mu\text{g/L}$ ).

In our sensitivity analysis, we used same linear dose-response model that was selected by the National Research Council in setting a reference dose for mercury exposure (NRC 2000). The Faroes researchers found that, for those children whose mothers had hair mercury concentrations  $<10 \mu\text{g/g}$ , a  $1 \mu\text{g/L}$  increase of cord blood mercury concentration was associated with adverse impacts on neurodevelopmental tests ranging from 3.95-8.33% of a standard deviation, or 0.59-1.24 IQ points (average=0.93 IQ points) (Jorgensen 2004). We also varied the cord/maternal blood mercury ratio from 1 to 1.7 in calculating IQ impact from the linear model as part of our sensitivity analysis. As an upper bound to our cost estimate using the logarithmic model, we calculated the economic cost assuming that children born to women with mercury concentrations 3.5-4.84  $\mu\text{g/L}$  suffer no loss in cognition, and that successive portions of the birth cohort experience losses of cognition of 1.21, 1.84, 2.55 and 5.13 IQ points, respectively. The lower bound estimate assumed that children born to women with mercury concentrations 4.84-5.8  $\mu\text{g/L}$  suffer no loss in cognition, and that successive portions of the birth cohort experience losses of cognition of 0.22, 0.48, and 1.39 IQ points.

As an upper bound to our cost estimate using the linear model, we calculated the economic cost assuming that children born to women with mercury concentrations 3.5-4.84  $\mu\text{g/L}$  suffer no loss in cognition, and that successive portions of the birth cohort experience losses of cognition of 3.01, 5.04, 7.84 and 24.43 IQ points, respectively. The lower bound estimate assumed that children born to women with mercury concentrations 4.84-5.8  $\mu\text{g/L}$  suffer no loss in cognition,

and that successive portions of the birth cohort experience losses of cognition of 0.56, 1.35 and 5.99 IQ points.

Calculation of Economic Costs of IQ Loss. To estimate the costs associated with the cognitive and behavioral consequences of mercury exposure, we relied on an economic forecasting model developed by Schwartz et al. and we applied this model to NHANES data on prevalence of mercury exposure in women of childbearing age (Schober et al. 2003; Schwartz et al. 1985). In this model, lead concentrations are assumed on the basis of work by Salkever to produce a dose-related decrement in IQ score. Those decrements in IQ are, in turn, associated with lower wages and diminished lifetime earning power. Salkever used three regression techniques to derive direct and indirect relationships amongst IQ, schooling, probability of workforce participation and earnings. He estimated a percentage in lost earnings per IQ point from the percent loss of earnings per  $\mu\text{g/dL}$  increase in blood lead level. Salkever found a .473 point decrement in lost lifetime earnings per  $\mu\text{g/dL}$  increase amongst men and a .806 point decrement per  $\mu\text{g/dL}$  increase amongst women. (Salkever 1995) Using Schwartz's estimate that 0.245 IQ points are lost per  $\mu\text{g/dL}$  increase in blood lead (Schwartz 1994), Salkever estimated a percentage loss in lifetime earnings per IQ point amongst men (1.931%) and women (3.225%). (Salkever 1995) We assume that this relationship remains linear across the population range of IQ.

Assuming an annual growth in productivity of 1% and applying a 3% real discount rate, the present value of lifetime expected earnings is \$1,032,002 for a boy born in 2000, and \$763,468 for a girl born in the same year (Max et al. 2002). The costs of the diminution in this earning power were calculated for the 2000 American birth cohort, using available data on the number of

male and female births in 2000 (U.S. CDC 2002a). We diminished our cost estimate by .69%, the infant mortality rate in 2000, to account for those children for whom methylmercury exposure is unlikely to result in diminished economic productivity. (U.S. CDC 2002b)

*American Sources of Mercury Emission.* Mercury emissions result from anthropogenic as well as from natural sources, and we chose to limit our analysis to methylmercury derived from anthropogenic sources. The United Nations Environment Programme recently estimated that anthropogenic uses account for 70% of the 5500 tons of mercury released into the earth's atmosphere worldwide (United Nations Environment Programme 2002). Therefore, to limit our analysis to anthropogenic mercury, we applied a 70% factor to convert the cost of lost economic productivity resulting from methylmercury exposure to the cost attributable to anthropogenic methylmercury exposure.

We next parsed out the proportion of anthropogenic methylmercury in fish that arises from American sources, and then isolated the subset of that proportion that is emitted by coal-fired electrical generating plants. In 1995, the most recent year for which federal data on the relative deposition of mercury from American and other global sources are available, 158 tons of mercury were emitted to the atmosphere by American anthropogenic sources. Fifty-two (33%) of those 158 tons were deposited in the lower forty-eight states, while the remaining two-thirds were added to the global reservoir (U.S. EPA 2004e). In that year, an additional thirty-five tons of mercury from the global reservoir were deposited in the United States. Therefore, a total of eighty-seven total tons of mercury were deposited in the United States in that year, of which 60% (52/87) were attributable to American anthropogenic sources (U.S. EPA 1997; U.S. EPA 1996).

This mercury would have been available to bioaccumulate in the marine and aquatic food chains and to enter American fresh- and saltwater fish.

Further complicating our calculations is the fact that not all of the fish sold in America is from American sources. Of the 10.4 billion pounds of edible fish supplied in the U.S. in 2002, 4.4 billion (42%) are imported from sources outside of the United States (National Marine Fisheries Service 2002). Since U.S. emissions account for 3% of global emissions (United Nations Environment Programme 2002; U.S. EPA 1996), we calculate that the mercury content of imported fish is 2% of American anthropogenic origin:

$$\begin{aligned} & 158 \text{ tons of American emissions-} \\ & 52 \text{ tons of American mercury deposited on American soil} \\ & = 106 \text{ tons of American mercury available to contaminate imported fish} \end{aligned}$$

---

$$\begin{aligned} & 5500 \text{ tons emitted globally} \\ & -87 \text{ tons deposited on American soil} \\ & = 5413 \text{ tons of mercury from all sources to contaminate imported fish} \\ & = 106/5413 = 2\% \text{ of mercury in imported fish of American origin} \end{aligned}$$

In the remaining 58% of fish consumed in the United States, we assume that 60% of the mercury content comes from American anthropogenic sources (U.S. EPA 1997; U.S. EPA 1996). We therefore applied a 36% factor (the weighted average of American sources of mercury content in fish, or  $0.6*0.58+0.02*0.42$ ) to specify the economic costs of anthropogenic methylmercury exposure attributable to American sources.

Modeling supported by the Electric Power Resource Institute (EPRI) estimates that 70% of the mercury deposited in the U.S. comes from foreign sources. This EPRI analysis also finds that U.S. sources are responsible for more than 60% of mercury deposition in the Boston–Washington, D.C. corridor. In one of the model’s selected receptor areas, Pines Lake, New Jersey, 80% of the deposition originated from U.S. sources, showing that regional deposition can be higher than the 60% number we use in this analysis (Seigneur et al. 2004). In our sensitivity analysis, we varied the factor used to convert the economic cost of anthropogenic methylmercury exposure to the economic cost attributable to American sources from 18% ( $0.3 \times 0.58 + 0.02 \times 0.42$ , using EPRI’s modeling) to 36% (using federal data on mercury deposition).

In 1999, the most recent year for which data on American mercury emissions are available, 48 (41%) of the 117 tons of mercury emissions from anthropogenic sources in the United States, were emitted by electric power generation facilities (U.S. EPA 2003a). To calculate the economic cost of methylmercury exposure attributable to these facilities, we applied an additional fraction of 41% in our analysis.

## **Results**

*Base-case analysis:* Each year in the United States, between 316,588 (7.8% of the annual birth cohort), and 637,233 babies, are born with cord blood mercury levels  $>5.8 \mu\text{g/L}$ . The lower bound estimate of 316,588 babies is based on the very conservative assumption that maternal and cord blood mercury concentrations are equal. But if the cord blood mercury concentration is on average 70% higher than the maternal blood mercury concentration, as suggested by recent

research, (Stern and Smith 2003), 637,233 babies, or 15.7% of the birth cohort, experience cord blood mercury levels  $>5.8 \mu\text{g/L}$ . Fetal blood mercury levels above  $5.8 \mu\text{g/L}$  are associated with small but significant loss of IQ. This decrement in IQ appears to be permanent and irreversible, and it adversely affects a significant portion of the annual birth cohort's economic productivity over a lifetime.

Using our base case assumptions (impact for women with total Hg  $>4.84 \mu\text{g/L}$ , cord/maternal Hg ratio=1.7, IQ impact 1.5 points per doubling), we calculated costs for the 405,881 children who suffer IQ decrements resulting from fetal methylmercury exposure. Under these assumptions, 89,293 children suffered a 0.76 decrement in IQ and another 113,647 experienced a 1.15 IQ point decrement. The 5% most highly exposed children in the 2000 birth cohort suffered subclinical losses in IQ in our model ranging from 1.60-3.21 points. While this diminution in intelligence is small in comparison with the loss of cognition that can result from other genetic and environmental processes, the loss resulting from methylmercury exposure produces a significant reduction in economic productivity over a lifetime. We estimate the aggregate cost of the loss in IQ that results from exposure of American children to methylmercury of anthropogenic origin to be \$8.7 billion (in 2000 dollars) annually (Table 1).

*Sensitivity Analysis:* We estimate that the cost of anthropogenic methylmercury exposure ranges from \$2.2 (impact only for the 316,588 children born to women with total Hg  $>5.8 \mu\text{g/L}$ , IQ impact 0.85 points per doubling) to \$13.9 billion (impact for the 405,881 women with total Hg  $>4.84 \mu\text{g/L}$ , IQ impact 2.4 points per doubling). Using the linear dose-response model that was selected by the National Research Council in recommending a reference dose for mercury

exposure (a model that predicts an average loss of 0.93 IQ points per 1  $\mu\text{g/L}$  increase in mercury concentration), (NAS 2000; Jorgensen et al. 2004) we find that the environmentally attributable cost of methylmercury exposure is \$32.9 billion, assuming a cord/maternal blood Hg ratio of 1.7. Employing a linear model and assuming that the true loss in IQ resulting from a 1  $\mu\text{g/L}$  increase in blood mercury ranges from 0.59 to 1.24 points, we find that the outer bounds of our estimate range from \$7.0 (impact only for women with total Hg  $>5.8 \mu\text{g/L}$ , IQ impact 0.59 points per  $\mu\text{g/L}$  increase, cord/maternal Hg ratio=1) to \$43.8 billion (impact for women with total Hg  $>4.84 \mu\text{g/L}$ , IQ impact 1.24 points per  $\mu\text{g/L}$  increase, cord/maternal Hg ratio=1.7). (Table 2)

*Sources of Costs:* After applying the 36% fraction to restrict our analysis to American anthropogenic sources, we estimate that the attributable cost of methylmercury exposure to the developing fetus from American anthropogenic sources is \$3.1 billion annually, using the logarithmic model developed by the Faroes researchers (Grandjean et al. 1999; Jorgensen et al. 2004) and assuming a 1.5 point IQ impact for each doubling of methylmercury exposure (Budtz-Jorgensen et al. 2002). Our sensitivity analysis, in which we also varied the attributable fraction for American sources from 18% (industry data sources) to 36% (federal data sources), suggests that the true cost of methylmercury exposure from American emissions ranges from \$0.4 to \$15.8 billion annually.

To focus specifically on the costs of fetal exposure to mercury released by American coal-fired power plants, we examined the impact of the 41% of U.S. anthropogenic emissions of mercury attributable to these facilities. We estimate that the attributable cost of methylmercury exposure from American electric generation facilities to the developing fetus is \$1.3 billion. Applying our

sensitivity analysis in this model, we find that the true cost of methylmercury exposure from electric generation facilities to the American birth cohort ranges from \$0.1 to \$6.5 billion/year (Figure 1). Again the major source of these costs is loss of earnings over a lifetime.

## **Discussion**

The major findings in this analysis are: (1) that exposure to methylmercury emitted to the atmosphere by American electric generation facilities causes lifelong loss of intelligence in hundreds of thousands of American babies born each year, and (2) that this loss of intelligence exacts a significant economic cost to American society, a cost that amounts to at least hundreds of million dollars each year. Moreover, these costs will recur each year with each new birth cohort so long as mercury emissions are not controlled. By contrast, the cost of installing stack filters to control atmospheric mercury emissions is a one-time expense. The high costs of *in utero* exposure to methylmercury are due principally to the lifelong consequences of irreversible injury to the developing brain. Similar lifelong neurobehavioral consequences have been observed following exposure of the developing brain to other environmental toxicants, including lead (Baghurst et al. 1987; Bellinger 2004; Dietrich et al. 1987; Opler et al. 2004; Wasserman et al. 2000), polychlorinated biphenyls (Jacobson and Jacobson 1996) and ethanol (Lupton et al. 2004).

Because the literature has presented a range of possible consequences for methylmercury toxicity, we have provided a range of possible public health and economic consequences. This range is meant to inform the choices that environmental and public health officials make in

protecting vulnerable populations from methylmercury exposure. Our range for the true economic costs of methylmercury toxicity to the developing brain does not capture the cost of exposures to the 231,352 children born to women in 2000 with blood mercury concentrations between 3.5 and 4.84  $\mu\text{g/L}$ . If the true cord blood ratio is 1.7 times the maternal blood concentration, as described in the most recent and extensive metaanalysis on the matter, (Stern and Smith 2003) these children are also born with cord blood mercury concentrations above the 5.8  $\mu\text{g/L}$  concentration at which adverse neurodevelopmental impact has been found. We chose not to include them in our analysis because other studies have found lower ratios and because we restricted ourselves in this analysis to the use of available, published prevalence data of maternal blood mercury concentrations. In our sensitivity analysis, we also selected low cord/maternal blood ratios so as to describe most accurately the range of values for the true cost of methylmercury exposure to the developing fetus.

Our analysis also fails to capture the cost of the cardiovascular impacts of mercury exposure (Grandjean et al. 2004), or the costs of mercury exposure to children in the first two years of postnatal life, when myelination is still continuing and the blood brain barrier remains vulnerable to penetration by methylmercury (Rodier 1995). We chose not to include these aspects of methylmercury toxicity in our range of estimates at this time, because there do not exist sufficient quantitative data to permit construction of a reliable model.

A limitation on our analysis is that it did not consider other societal costs beyond decreased lifetime earnings that may result from exposure of the developing brain to methylmercury. For example, if the value of a child's social productivity is approximately \$4-9 million, as suggested

by studies of willingness to pay (WTP) estimates of a life (Viscusi et al. 2004), then by the WTP methodology the true cost of methylmercury toxicity may be much higher than our estimate. We also chose not to include other non-cognitive impacts. Lead, for example, has been associated with criminality and antisocial behavior (Dietrich et al 2001; Needleman et al 1996, 2002; Nevin 2000; Stretesky and Lynch 2001). However, because these behaviors have not been described as yet for methylmercury, we chose not to include such costs in our estimate.

Some will argue that our range of costs fails to incorporate the role of confounding factors in quantifying the economic consequences of methylmercury exposure. It is true that efforts to delineate the potential synergistic role of methylmercury and other chemicals in mediating neurocognitive and other effects are bedeviled by lack of knowledge about possible interactions and synergies among chemicals or between chemicals and other environmental hazards, even though the environment of a child includes mixtures of chemical and biological toxicants. Only a study of the magnitude of the National Children's Study will facilitate simultaneous examination of the effects of multiple chemical exposures, of interactions among them, and of interactions among biologic, chemical, behavioral, and social factors (Trasande and Landrigan 2004). However, we note that loss of cognition resulting from methylmercury exposure in the Faroe Islands study remained evident when blood levels of polychlorinated biphenyls, which are known fetal neurotoxicants (Jacobson and Jacobson 1996), were included in the analysis (Budtz-Jorgensen et al. 2002; Steuerwald et al., 2000).

We note the EPA's recent success in minimizing mercury emissions from medical waste (U.S. EPA 2004f) and municipal incinerators (U.S. EPA 2004g; U.S. EPA 2004h), actions which

resulted in a decrease in total mercury emissions by at least 80 tons per year from 1990 to 1999 (U.S. EPA 2003b). Though data are not available on blood mercury concentrations over the past decade that followed from those actions, the impact of these reductions is likely to have been substantial.

Some commentators have used data from the Seychelles study to argue that methylmercury is not toxic to the fetus at low concentrations and to suggest that fear of mercury exposure is needlessly preventing women from ingesting fish, and thus denying them access to beneficial long-chain polyunsaturated fatty acids (LCPUFAs), especially docosahexaenoic acid (DHA). We do not dispute that DHA and other LCPUFAs are important for optimal development of the fetal visual and nervous systems (Innis 1991). The human fetus has a limited ability to synthesize DHA's precursor,  $\alpha$ -linolenic acid, and therefore it must be largely supplied from maternal sources (Carnielli et al. 1996; Larque et al 2002; Szitanyi et al. 1999). We also note a report that associated an average monthly decline in fish consumption of 1.4 servings among Massachusetts women with an FDA advisory on the health risks of mercury (Oken et al. 2003). Nonetheless, the American Heart Association, a strong advocate for the cardioprotective effects of LCPUFAs, recommends the avoidance of potentially contaminated fish in children and pregnant and lactating women (Kris-Etherton 2002). Fish advisories should not recommend that consumers abstain from fish, but they should assist in choosing the best kinds of fish to eat. Lists of fish that are safe and unsafe from the perspective of mercury exposure have been published and made widely available to consumers (U.S. EPA 2004i).

Early reports of disease and dysfunction of environmental origin in children have on repeated occasions failed to produce proactive response to protect children. The long history of lead use in the U.S. provides a chilling reminder of the consequences of failure to act on early evidence of harm. It is important that we not repeat this sequence with mercury. Within the last century, as a result of increased industrial activity, mercury emissions worldwide have increased two- to five-fold, and anthropogenic emissions now surpass emissions from natural sources. (Nriagu 1989)

The data from this analysis reinforce the results of recent epidemiological studies and indicate an urgent need on economic grounds for regulatory intervention at the federal level to minimize mercury emissions. Our analysis captures the cost of methylmercury exposure for only one year's birth cohort, but the cost of mercury exposure will continue to accrue in each succeeding year if power plants fail to install flue gas filters (U.S. Department of Energy 2004) or to implement other technologies to reduce mercury emissions. The cost savings from reducing mercury exposure now will provide savings in improved productivity and enhanced national security for generations to come.

## References

Amin-Zaki L, Elhassani S, Majeed MA, Clarkson TW, Doherty RA, Greenwood M. 1974. Intrauterine methylmercury poisoning in Iraq. *Pediatrics* 54:587-595.

Amin-Zaki L, Elhassani S, Majeed MA, Clarkson TW, Greenwood M, Doherty RA. 1979. Prenatal methylmercury poisoning. Clinical observations over five years. *Am J Dis Child* 133:172-177.

Baghurst PA, Robertson EF, McMichael AJ, Vimpani GV, Wigg NR, Roberts RR. 1987. The Port Pirie Cohort Study: lead effects on pregnancy outcome and early childhood development. *Neurotoxicology*. 8(3):395-401.

Bakir F, Damluji SF, Amin-Zaki L, Murtadha M, Khalidi A, al-Rawi NY, et al. 1973. Methylmercury poisoning in Iraq. *Science*. 181:230-241.

Bellinger DC. Lead. *Pediatrics*. 2004 Apr;113(4 Suppl):1016-22.

Budtz-Jørgensen E, Grandjean P, Jørgensen PJ, Weihe P, Keiding N. 2004. Association between mercury concentrations in blood and hair in methylmercury-exposed subjects at different ages. *Environmental Research*, in press. Available at <http://www.chef-project.dk>. (Accessed on 7 June 2004)

Budtz-Jorgensen E, Keiding N, Grandjean P, Weihe P. 2002. Estimation of health effects of prenatal methylmercury exposure using structural equation models. *Environ Health* 1(1):2.

Campbell D, Gonzales M, Sullivan JB Jr. 1992. Mercury. In: Sullivan JB Jr, Krieger GR, eds. *Hazardous Materials Toxicology—Clinical Principles of Environmental Health*. Baltimore, MD: Williams and Wilkins; 1992: 824-833.

Carnielli VP, Wattimena DJ, Luijendijk IH, Boerlage A, Degenhart HJ, Sauer PJ. 1996. The very low birth weight premature infant is capable of synthesizing arachidonic and docosahexaenoic acids from linoleic and linolenic acids. *Pediatric Res* 40:169–74.

Dietrich KN, Krafft KM, Bornschein RL, Hammond PB, Berger O, Succop PA, et al. 1987. Low-level fetal lead exposure effect on neurobehavioral development in early infancy. *Pediatrics* 80(5):721-730.

Dietrich KN, Ris MD, Succop PA, Berger OG, Bornschein RL. Early exposure to lead and juvenile delinquency. *Neurotoxicol Teratol* 23:511-518 (2001).

Dietz R, Riget F, Cleemann M, Aarkrog A, Johansen P, Hansen JC. 2000. Comparison of contaminants from different trophic concentrations and ecosystems. *Science of the Total Environment* 245 (1-3):221-223.

Fahs MC, Markowitz SB, Fischer E, Shapiro J, Landrigan PJ. 1989. Health costs of occupational disease in New York State. *Am J Ind Med* 16:437-449.

Franzblau A. Mercury. p. 756-759. In: Rosenstock L, Cullen M. 1994. *Textbook of Occupational and Environmental Medicine*. Philadelphia: W.B. Saunders Company.

Gilmour CC, Riedel GS. 2000. A survey of size-specific mercury concentrations in game fish from Maryland fresh and estuarine waters. *Archives of Environmental Contamination and Toxicology* 39 (1):53-59.

Goldman LR, Shannon MW; American Academy of Pediatrics: Committee on Environmental Health. 2001. Technical report: mercury in the environment: implications for pediatricians. *Pediatrics*. 108(1):197-205.

Grandjean P, Weihe P, White RF, Debes F, Araki S, Yokoyama K, et al. 1997. Cognitive Deficit in 7-Year-Old Children with Prenatal Exposure to Methylmercury. *Neurotoxicol Teratol* 19(6):417-428.

Grandjean P, Murata K, Budtz-Jorgensen E, Weihe P. 2004. Cardiac autonomic activity in methylmercury neurotoxicity: 14-year follow-up of a Faroese birth cohort. *J Pediatr* 144(2):169-176.

- Grandjean P, Budtz-Jorgensen E, White RF, Jorgensen PJ, Weihe P, Debes F, et al. 1999. Methylmercury exposure biomarkers as indicators of neurotoxicity in children age 7 years. *Am J Epidemiol* 150:301-305.
- Guimaraes JRD, Ikingura J, Akagi H. 2000. Methyl mercury production and distribution in river water-sediment systems investigated through radiochemical techniques. *Water, Air, and Soil Pollution* 124 (1-2):113-124.
- Harada Y. Congenital (or fetal) Minamata disease. 1968. In: Study Group of Minamata Disease, eds. *Minamata Disease*. Kumamoto, Japan: Kumamoto University. 93-118.
- Institute of Medicine. 1981. *Costs of Environment-Related Health Effects: A Plan for Continuing Study*. Washington, DC:National Academy Press.
- Innis SM. Essential fatty acids in growth and development. 1991. *Prog Lipid Res* 30:39–103.
- Jacobson JL, Jacobson SW. 1996. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* 335:783-789.
- Jorgensen EB, Weihe P, Grandjean P. 2004. Adverse mercury effects in seven year old children as expressed as loss in IQ. Available at [www.chef-project.dk](http://www.chef-project.dk) (Accessed 15 May 2004.)

Kjellstrom T, Kennedy P, Wallis S, Mantell C. 1986. Physical and Mental Development of Children With Prenatal Exposure to Mercury From Fish. Stage I: Preliminary Tests at Age 4. Solna, Sweden: National Swedish Environmental Protection Board (Report 3080)

Kjellstrom T, Kennedy P, Wallis S, Stewart A, Friberg L, Lind B et al. 1989. Physical and Mental Development of Children With Prenatal Exposure to Mercury From Fish. Stage II: Interviews and Psychological Tests at Age 6. Solna, Sweden: National Swedish Environmental Protection Board (Report 3642)

Kris-Etherton PM, Harris WS, Appel LJ, for the Nutrition Committee. 2002. Fish Consumption, Fish Oil, Omega-3 Fatty Acids, and Cardiovascular Disease. *Circulation* 106:2747-2757.

Landrigan PJ, Schechter CB, Lipton JM, Fahs MC, Schwartz J. 2002. Environmental pollutants and disease in American children: estimates of morbidity, mortality and costs for lead poisoning, asthma, cancer and developmental disabilities. *Environ Health Perspect* 110:721-728.

Landrigan PJ, Goldman L. 2003. Prenatal methylmercury exposure in the Seychelles. *Lancet* 362(9384):666.

Larque E, Demmelmair H, Koletzko B. 2002. Perinatal supply and metabolism of long-chain polyunsaturated fatty acids: importance for the early development of the nervous system. *Ann NY Acad Sci* 967:299-310.

Lupton C, Burd L, Harwood R. 2004. Cost of fetal alcohol spectrum disorders. *Am J Med Genet* 127C(1):42-50.

Leigh JP, Markowitz S, Fahs M, Shin C, Landrigan PJ. 1997. Costs of occupational injuries and illnesses. *Arch Intern Med* 157:1557-1568.

Mahaffey KR, Clickner RP, Boudrow CC. 2004. Blood Organic Mercury and Dietary Mercury Intake: National Health and Examination Survey, 1999 and 2000. *Environmental Health Perspect* 112(5):562-570.

Mason RP, Reinfelder JR, Morel FMM. 1995. Bioaccumulation of mercury and methylmercury. *Water, Air, and Soil Pollution* 80:915-921.

Max W, Rice DP, Sung H-Y, Michel M. 2002. *Valuing Human Life: Estimating the Present Value of Lifetime Earnings*, 2000. San Francisco: Institute for Health & Aging.

Myers GJ, Davidson PW, Cox C, Shamlaye CF, Palumbo D, Cernichiari E et al. 2003. Prenatal methylmercury exposure from the ocean fish consumption in the Seychelles child development study. *Lancet* 361:1686-1692.

Murata K, Weihe P, Budtz-Jorgensen E, Jorgensen PJ, Grandjean P. 2004. Delayed brainstem auditory evoked potential latencies in 14-year-old children exposed to methylmercury. *J Pediatr* 144(2):177-183.

National Research Council. 2000. Toxicological Effects of Methylmercury. Washington, DC: National Academy Press.

National Marine Fisheries Service, National Oceanic and Atmospheric Administration. 2004. Fisheries of the United States – 2002. Available at <http://www.st.nmfs.gov/st1/fus/current/2002-fus.pdf> (Accessed 24 May 2004).

Needleman HL, Riess JA, Tobin MJ, Biesecker GE, Greenhouse JB Bone lead levels and delinquent behavior. JAMA 275:363-369 (1996).

Needleman HL, McFarland C, Ness RB, Fienberg SE, Tobin MJ. Bone lead levels in adjudicated delinquency: a case-control study. Neurotoxicol. Teratol. 24:711–17 (2002).

Neumann RM, Ward S.M. 1999. Bioaccumulation and biomagnification of mercury in two warmwater fish communities. Journal of Freshwater Ecology 14 (4):487-498.

Nevin R. How lead exposure relates to temporal changes in IQ, violent crime, and unwed pregnancy. Environ Res 83:1–22 (2000).

Nriagu, JO. 1989. A global assessment of natural sources of atmospheric trace metals. Nature 338: 47-49.

Oken E, Kleinman KP, Berland WE, Simon SR, Rich-Edwards JW, Gillman MW. 2003.

Decline in fish consumption among pregnant women after a national mercury advisory. *Obstet Gynecol* 102(2):346-351.

Opler MG, Brown AS, Graziano J, Desai M, Zheng W, Schaefer C, et al 2004. Prenatal lead exposure, delta-aminolevulinic Acid, and schizophrenia. *Environ Health Perspect* 112(5):548-552.

Rice DC, Schoeny R, Mahaffey K. 2003. Methods and rationale for a reference dose for methylmercury by the U.S. EPA. *Risk Anal* 23(1):107-115.

Rodier PM. 1995. Developing brain as a target of toxicity. *Environ Health Perspect* 103(suppl 6):S73-S76.

Salkever DS. 1995. Updated estimates of earnings benefits from reduced exposure of children to environmental lead. *Environ Res* 70:1-6.

Schwartz J, Pitcher H, Levin R, Ostro B, Nichols AL. 1985. *Costs and Benefits of Reducing Lead in Gasoline: Final Regulatory Impact Analysis*. EPA-230/05-85/006. Washington, DC:U.S. Environmental Protection Agency.

Schwartz J. 1994. Societal Benefits of Reducing Lead Exposure. *Environmental Research* 55:105-124.

Schober SE, Sinks TH, Jones RL, Bolger PM, McDowell M, Osterloh J, et al. 2003. Blood mercury concentrations in US children and women of childbearing age, 1999-2000. *JAMA* 289:1667-1674.

Seigneur C, Vijayaraghavan K, Lohman K, Karamchandani P and Scott C. 2004. Global Source Attributions for Mercury Deposition in the United States. *Environ Sci Technol.* 38(2):555-569.

Smith KR, Corvalin CF, Kjellstrom T. 1999. How much global ill health is attributable to environmental factors? *Epidemiology* 10:573-584.

Stern AH. 2005. A revised probabilistic estimate of the maternal methyl mercury intake dose corresponding to a measured cord blood mercury concentration. *Environ Health Perspect* 113:155-163.

Stern AH, Smith AE. 2003. An assessment of the cord blood-maternal blood methylmercury ratio: implications for risk assessment. *Environ Health Perspect* 111:1465-1470.

Steuerwald U, Weihe P, Jorgensen PJ, Bjerve K, Brock J, Heinzow B, et al. 2000. Maternal seafood diet, methylmercury exposure and neonatal neurological function. *J Pediatr* 136: 599-605.

Stretesky PB, Lynch MJ. The relationship between lead exposure and homicide. *Pediatr. Adolesc. Med.* 155:579–82 (2001).

Szitanyi P, Koletzko B, Mydlilova A, Demmelhair H. 1999. Metabolism of <sup>13</sup>C-labeled linoleic acid in newborn infants during the first week of life. *Pediatric Res* 45:669–73.

Trasande L, Landrigan PJ. The National Children's Study: A Critical National Investment. *Environmental Health Perspectives* 2004;112(14):A789-790.

United Nations Environmental Programme. 2002. Global Mercury Assessment Report. Available at <http://www.chem.unep.ch/mercury/Report/GMA-report-TOC.htm> (Accessed 17 May 2004).

U.S. Centers for Disease Control. 2002a. National Vital Statistics System. Births: Final Data for 2000. *National Vital Statistics Report* 50(5):1-102.

U.S. Centers for Disease Control. 2002b. National Vital Statistics System. Infant Mortality Statistics from the 2000 Period Linked Birth/Infant Death Data Set. *National Vital Statistics Report* 50(12):1-27. Available at [http://www.cdc.gov/nchs/data/nvsr/nvsr50/nvsr50\\_12.pdf](http://www.cdc.gov/nchs/data/nvsr/nvsr50/nvsr50_12.pdf) [Accessed 1 November 2004.]

U.S. Department of Energy, Office of Fossil Energy. 2004. Mercury Controls. Available at [http://www.fossil.energy.gov/programs/powersystems/pollutioncontrols/overview\\_mercurycontrols.shtml](http://www.fossil.energy.gov/programs/powersystems/pollutioncontrols/overview_mercurycontrols.shtml) (Accessed 18 May 2004).

U.S. Environmental Protection Agency, Technology Transfer Network (TTN), Clearinghouse for Inventories and Emissions Factors. 2003a. National Emissions Inventories for Hazardous Air Pollutants, 1999. Version 3, July 2003. Available at <http://www.epa.gov/ttn/chief> (Accessed 18 May 2004).

U.S. Environmental Protection Agency, Technology Transfer Network (TTN), Clearinghouse for Inventories and Emissions Factors. 2003b. National Toxics Inventory, 1990, Version 0302, October 2003. Available at <http://www.epa.gov/ttn/chief> (Accessed 18 May 2004).

U.S. Environmental Protection Agency. 2004a. EPA Proposes Options for Significantly Reducing Mercury Emissions from Electric Utilities. Available at [http://www.epa.gov/mercury/hg\\_factsheet1\\_29\\_04.pdf](http://www.epa.gov/mercury/hg_factsheet1_29_04.pdf) (Accessed May 7, 2004).

U.S. Environmental Protection Agency. 2004b. Clean Air Act. Available at [http://www.epa.gov/oar/oaq\\_caa.html](http://www.epa.gov/oar/oaq_caa.html) (Accessed 17 May 2004).

U.S. Environmental Protection Agency. 2004c. Section B: Human and Environmental Benefits. 2002 Technical Support Package for Clear Skies. Available at [http://www.epa.gov/clearskies/tech\\_sectionb.pdf](http://www.epa.gov/clearskies/tech_sectionb.pdf) (Accessed 17 May 2004).

U.S. Environmental Protection Agency. 2004d. Utility Mercury Reductions Rule: Basic Information. Available at <http://www.epa.gov/air/mercuryrule/basic.htm> (Accessed 1 June 2004).

U.S. Environmental Protection Agency. 2004e. Mercury study report to Congress: Overview. Available at <http://www.epa.gov/oar/mercover.html> [Accessed 1 November 2004.]

U.S. Environmental Protection Agency. 2004f. Hospital/Medical/Infectious Waste Incinerators. <http://www.epa.gov/ttnatw01/129/hmiwi/rihmiwi.html> (Accessed on 18 May 2004).

U.S. Environmental Protection Agency. 2004g. Rule and Implementation Information for Large Municipal Waste Combustors. <http://www.epa.gov/ttn/atw/129/mwc/rimwc.html> (Accessed on 18 May 2004).

U.S. Environmental Protection Agency. 2004h. Rule and Implementation Information for Small Municipal Waste Combustors. Available at <http://www.epa.gov/ttn/atw/129/mwc/rimwc2.html> (Accessed on 18 May 2004).

U.S. Environmental Protection Agency. 2004i. Links to Local Fish Advisory Programs. Available at <http://epa.gov/waterscience/fish/states.htm> [Accessed 9 November 2004.]

U.S. Environmental Protection Agency. 1996. Mercury study report to Congress. Volume II: An inventory of anthropogenic mercury emissions in the United States. EPA452/R-96-001b. Washington, DC: U.S. Environmental Protection Agency.

U.S. Environmental Protection Agency. 1997. Locating and estimating air emissions from sources of mercury and mercury compounds. Office of Air Quality Planning and Standards and Office of Air and Radiation. Research Triangle Park, NC. EPA-454/R-97-012. Washington, DC: U.S. Environmental Protection Agency.

Viscusi WK, Aldy JE. 2004. The Value of a Statistical Life: A Critical Review of Market Estimates Throughout the World. NBER Working Paper Series, Working Paper 9487. Available at <http://www.nber.org/papers/w9487> (Accessed on 2 July 2004).

Wasserman GA, Liu X, Popovac D, Factor-Litvak P, Kline J, Wateraux C, et al. 2000. The Yugoslavia Prospective Lead Study: contributions of prenatal and postnatal lead exposure to early intelligence. *Neurotoxicol Teratol* Nov-Dec;22(6):811-818.

Table 1. Cost of Anthropogenic Mercury Exposure Using Logarithmic Model

Segment of Population	90-92.1%ile Hg	92.2-94.9%ile	95-99.3%ile	99.4%ile and up
Range of maternal total Hg concentration	4.84-5.8 µg/L	5.8-7.13 µg/L	7.13-15.0 µg/L	>15.0 µg/L
Assumed maternal total Hg concentration	4.84	5.8	7.13	15
No effect concentration (maternal total Hg)	3.41	3.41	3.41	3.41
IQ points lost at assumed concentration	0.76	1.15	1.60	3.21
Loss of 1 IQ point=	2.39% decrease in lifetime earnings			
For boys, lifetime earnings	\$1,032,002			
For girls, lifetime earnings	\$763,468			
Number of boys in birth cohort affected	45693	58155	91387	12462
Number of girls in birth cohort affected	43601	55492	87201	11891
Lost income	\$1.1 billion	\$2.0 billion	\$4.4 billion	\$1.2 billion
Total Cost	\$8.7 billion in each year's birth cohort			

Assumptions: Environmentally Attributable Fraction =70%, Main Consequence=Loss of IQ over lifetime

Table 2. Sensitivity Analysis, Cost of Anthropogenic Methylmercury Exposure

	Base case cost estimate (range of estimates based on range of possible IQ decrement/increase cord blood Hg)
Children born to women with Hg >4.84 µg/L, effect >3.5 µg/L	
Logarithmic Model	\$8.7 billion (\$4.9-13.9 billion)
Linear Model, Cord/Maternal Hg ratio=1.7	\$32.9 billion (\$20.9-\$43.8 billion)
Linear Model, Cord/Maternal Hg ratio=1	\$19.3 billion (\$12.3-\$25.8 billion)
	Base case cost estimate (range of estimates based on range of possible IQ decrement/increase cord blood Hg)
Children born to women with >5.8 µg/L, effect >4.84 µg/L	
Logarithmic Model	\$3.9 billion (\$2.2-\$6.3 billion)
Linear Model, Cord/Maternal Hg ratio=1.7	\$18.7 billion (\$11.9-\$24.9 billion)
Linear Model, Cord/Maternal Hg ratio=1	\$11.0 billion (\$7.0-\$14.6 billion)
	Range of Estimates
Logarithmic Model	\$2.2-\$13.9 billion
Linear Model	\$7.0-\$43.8 billion
Assumptions: Environmentally Attributable Fraction =70%, Main Consequence=Loss of IQ over lifetime	




Figure 1. Portions of Cost of Methylmercury Exposure Attributed to Sources  
 Assumptions: 18-36% attributable to American sources; 41% of American emissions attributable to American power plants